

# Talking With Your Doctor About LIVMARLI



Making treatment decisions for Alagille syndrome can be difficult. Speaking with your doctor can help make things easier. If you or your child are considering a change in your current treatment plan or are ready to start treatment, you may want to talk to your doctor about LIVMARLI.

LIVMARLI is a once-daily oral medicine for cholestatic pruritus (itch) in patients with Alagille syndrome who are 1 year of age and older.

Use the information below, as well as the Patient Information on LIVMARLI.com, to get the conversation started with your doctor about whether LIVMARLI may be right for you or your child.

## WHAT TO SHARE WITH YOUR DOCTOR



How often does itching impact daily life?

- Never       Once or twice per month       Once or twice per week       Many times per week       Every day



How does itching impact your child's and/or your family's daily life?

---

---

---

---

---

---



How much relief do you or your child get from your current itch treatment?

- A lot of relief       Some relief       No relief at all       I am not currently treating my itch



I hope a treatment for Alagille syndrome can help with:

---

## WHAT TO ASK YOUR DOCTOR

### About LIVMARLI

- Could LIVMARLI be an option for me or my child?
- How does LIVMARLI work?
- How long will it take for LIVMARLI to work?
- How will I know if LIVMARLI is working?



### Dosing and Safety

- How is LIVMARLI taken?
- What are the possible side effects of LIVMARLI? How can they be managed?
- Can LIVMARLI be taken if I or my child have other health conditions or take other medicines?

### Support

- How do I get LIVMARLI?
- What support is available to help me or my child throughout treatment with LIVMARLI?

Visit [www.LIVMARLI.com](http://www.LIVMARLI.com) for more information and support.

## INDICATION

LIVMARLI is a prescription medicine used to treat cholestatic pruritus (itch) in patients with Alagille syndrome 1 year of age and older. It is not known if LIVMARLI is safe and effective in children under 1 year of age or in adults 65 years and older.

## IMPORTANT SAFETY INFORMATION

### What are the possible side effects of LIVMARLI?

LIVMARLI can cause serious side effects, including:

**Changes in liver tests.** Changes in certain liver tests are common in patients with Alagille syndrome but may worsen during treatment with LIVMARLI. These changes may be a sign of liver injury and can be serious. Your health care provider should do blood tests before starting and during treatment to check your liver function. Tell your health care provider right away if you get any signs or symptoms of liver problems, including:

- nausea or vomiting
- skin or the white part of the eye turns yellow
- dark or brown urine
- pain on the right side of the stomach (abdomen)
- loss of appetite

**Stomach and intestinal (gastrointestinal) problems.** LIVMARLI can cause stomach and intestinal problems, including diarrhea, stomach pain, and vomiting during treatment. Tell your health care provider right away if you have any of these symptoms more often or more severely than normal for you.

A condition called **Fat Soluble Vitamin (FSV) Deficiency** caused by low levels of certain vitamins (vitamins A, D, E, and K) stored in body fat. FSV deficiency is common in patients with Alagille syndrome but may worsen during treatment. Your health care provider should do blood tests before starting and during treatment.

Other common side effects reported during treatment were bone fractures and gastrointestinal bleeding.

Tell your health care provider about all medicines that you take. LIVMARLI may affect the way some other medicines work, and some other medicines may affect the way LIVMARLI works. If you take a medicine that lowers cholesterol by binding bile acids, such as cholestyramine, colestevlam, or colestipol, take it at least 4 hours before or 4 hours after you take LIVMARLI.

LIVMARLI is taken by mouth, 1 time each day, 30 minutes before the first meal of the day. Be sure to use the provided oral dosing dispenser to accurately measure the dose of medicine.

These are not all of the possible side effects of LIVMARLI. Call your health care provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see full [Patient Information](#) for LIVMARLI.

